

Figure 1-Supplementary  
[Click here to download high resolution image](#)




















<u>ANIMAL GROUPS</u>		
<u>Group 1</u>	<u>Group 2</u>	<u>Group 3</u>
 <b>1-S-1</b>	 <b>1-S-2</b>	 <b>1-S-3 (a)</b>  <b>1-S-3 (b)</b>
<i>Normal histological structure</i>	<i>Nuclear pyknosis, degeneration and vacuolization with cytoplasmic oedema in neurons</i>	<i>(a) Nuclear pyknosis and degeneration in some neurons. (b) focal gliosis</i>
<u>Group 4</u>	<u>Group 5</u>	<u>Group 6</u>
 <b>1-S-4</b>	 <b>1-S-5</b>	 <b>1-S-6</b>
<i>The cerebral cortex shows nuclear pyknosis and degeneration in the neurons</i>	<i>Normal histological structure</i>	<i>Normal histological structure</i>



Figure 2-Supplementary  
[Click here to download high resolution image](#)

<u>ANIMAL GROUPS</u>		
<u>Group 1</u>	<u>Group 2</u>	<u>Group 3</u>
 <b>2-S-1 (a)</b>	 <b>2-S-2 (a)</b>	 <b>2-S-3 (a)</b>
 <b>2-S-1 (b)</b>	 <b>2-S-2 (b)</b>	 <b>2-S-3 (b)</b>
<i>No histopathological alterations are recorded with normal hippocampus (a) subiculum (b) fascia dentate</i>	<i>Nuclear degeneration is detected in the neurons of the both (a) subiculum (b) fascia dentate</i>	<i>Nuclear pyknosis and degeneration are identified in some (a) subiculum and (b) fascia dentate</i>
<u>Group 4</u>	<u>Group 5</u>	<u>Group 6</u>
 <b>2-S-4 (a)</b>	 <b>2-S-5 (a)</b>	 <b>2-S-6 (a)</b>
 <b>2-S-4 (b)</b>	 <b>2-S-5 (b)</b>	 <b>2-S-6 (b)</b>
<i>Normal histological structure is recorded in (a) subiculum and (b) fascia dentate</i>	<i>No histopathological alterations are identified in (a) subiculum and (b) fascia dentate</i>	<i>No histopathological alterations are identified</i>